

Non-Compact AdS_5 Universe with Parallel Positive Tension 3-Branes

Tianjun Li

Department of Physics, University of Wisconsin, Madison, WI 53706, U. S. A.

Abstract

We construct the general model with only parallel positive tension 3-branes on $M^4 \times R^1$ and $M^4 \times R^1/Z_2$. The cosmology constant is sectional constant along the fifth dimension. In this general scenario, the 5-dimensional GUT scale on each brane can be indentified as the 5-dimensional Planck scale, but, the 4-dimensional Planck scale is generated from the low 4-dimensional GUT scale exponentially in our world. We also give two simple models to show explicitly how to solve the gauge hierarchy problem.

November 1999

1 Introduction

Experiments at LEP and Tevatron have given the strong support to the Standard Model of the Strong and electroweak interactions. However, the Standard Model has some unattractive features which may imply the new physics beyond the Standard Model. One of these problems is that the gauge forces and the gravitational force are not unified. Another is the gauge hierarchy problem between the weak scale and the 4-dimensional Planck scale. Previously, two solutions to the gauge hierarchy problem have been proposed: one is the idea of the technicolor and compositeness which lacks calculability, and the other is the idea of supersymmetry.

More than one year ago, it was suggested that the large compactified extra dimensions may also be the solution to the gauge hierarchy problem [1]. Moreover, half year ago, Randall and Sundrum [2] proposed another scenario that the extra dimension is an orbifold, and the size of the extra dimension is not large but the 4-dimensional mass scale in the standard model is suppressed by an exponential factor from 5-dimensional mass scale. In addition, they suggested that the fifth dimension might be infinity [3], and there may exist only one brane with positive tension at the origin, but, there exists the gauge hierarchy problem. The remarkable aspect of the second scenario is that it gives rise to a localized graviton field. Combining those results, Lykken and Randall obtained the following physical picture [4]: the graviton is localized on Planck brane, we live on a brane separated from the Planck brane about 30 Planck lengths along the fifth dimension. On our brane, the mass scale in the Standard Model is suppressed exponentially, which gives the low energy scale. Furthermore, we generalized those scenarios and obtained the scenario with the following property [16]: the 4-dimensional Planck scale is generated from the low 5-dimensional Planck scale by an exponential hierarchy, and the mass scale in the Standard model, which is contained in the observable brane, is not rescaled. In short, recently, this kind of compactification or similar idea has attracted a lot of attentions [5-27]. By the way, supergravity domain walls was discussed previously in the 4-dimensional space-time [28].

In the model building, almost all the models [2, 4, 11, 16, 19, 26, 27] which can solve the gauge hierarchy problem contain the branes with the negative brane tension except Lykken-Randall model and Oda model ¹. However, mathematically speaking, Lykken-Randall model [4] may not be strict because they neglected the effect from the observable brane tension which they assumed might be very small, and in Oda model [26], the transformation is not 1-1 and may not be topological invariant. We want to consider the model with only positive tension branes, because, as we know, an object with negative tension can not be stable, although we might not need to worry about it if the anti-brane is orientable ². In addition, there are positive energy objects, namely D-branes and NS-branes, that are well understood and on which

¹ I do not consider the models with brane intersections/junctions here [6, 25], and many brane junctions will be considered elsewhere [18].

² I would like to thank J. X. Lu for pointing out this to me.

gauge fields and matter fields can be localized so that the Standard Model fields can be placed there.

Before our discussion, we would like to explain our assumption and notation which are similar to those in [16]. We assume that all the gauge forces are unified on each 3-brane if there exist gauge forces. The 5-dimensional GUT scale on i -th 3-brane $M_{GUT}^{(i)}$ and the 5-dimensional Planck scale M_X are defined as the GUT scale and Planck scale in the 5-dimensional fundamental metric, respectively. The 4-dimensional GUT scale on i -th 3-brane $M_{GUT}^{(i)}$ and the 4-dimensional Planck scale M_{pl} are defined as the GUT scale and Planck scale in the 4-dimensional Minkowski Metric ($\eta_{\mu\nu}$). In order to avoid the gauge hierarchy problem between the weak scale and the 4-dimensional GUT scale M_{GUT} on the observable brane which includes our world, we assume the low energy unification³. The key ansatz is that the 5-dimensional GUT scale on each brane is equal to the 5-dimensional Planck scale.

In this paper, we construct the general model with only parallel positive tension 3-branes. If the fifth dimension is compact, the sum of the brane tensions is zero which is the topological invariant of this kind of models [17]. Therefore, we have to go to the non-compact fifth dimension case. From differential topology/manifold, up to diffeomorphic, there is only one connected non-compact 1-dimensional manifold: R^1 , and there is only one connected non-compact 1-dimensional manifold with boundary: H_1 or R^1/Z_2 (the equivalence class is $y \sim -y$). Therefore, the space-time we consider is $M^4 \times R^1$ and $M^4 \times R^1/Z_2$ where M^4 is the 4-dimensional Minkowski space-time. The parallel positive tension 3-branes are located along the fifth dimension. We construct the most general model with above properties, calculate the 4-dimensional Planck scale and GUT scale on each brane. It seems to us the gauge hierarchy problem can be solved easily in this scenario: the 5-dimensional GUT scale on each brane can be indentified as the 5-dimensional Planck scale, but, the 4-dimensional Planck scale is generated from the low 4-dimensional GUT scale exponentially in our world. The cosmology constant is sectional constant along the fifth dimension, so, for any point in $M^4 \times R^1$ and $M^4 \times R^1/Z_2$, which is not belong to any brane and the section with zero cosmology constant, there is a neighborhood which is diffeomorphic to (or a slice of) AdS_5 space. Furthermore, We also give two simple models which are subsets of the model with more branes: general two branes case, three branes with Z_2 symmetry. The gauge hierarchy problem is solved explicitly in those models. One new feature is that, sometimes, the 5-dimensional Planck scale might be very large because of the exponential factor.

³We will not explain why M_{GUT} can be low energy scale here, but, it is possible if one considers additional particles which change the RGE running. Of course, proton decay might be the problem, but we do not discuss this here. These subjects were discussed in [19, 29].

2 General Solution and Model

First, we consider the fifth dimension is R^1 . Assuming we have $l + m + 1$ parallel positive tension 3-branes, and their fifth coordinastes are: $-\infty < y_{-l} < y_{-l+1} < \dots < y_{-1} < y_0 < y_1 < \dots < y_{m-1} < y_m < +\infty$. The 5-dimensional metric in these branes are:

$$g_{\mu\nu}^{(i)}(x^\mu) \equiv G_{\mu\nu}(x^\mu, y = y_i) , \quad (1)$$

where G_{AB} is the five-dimensional metric, and $A, B = \mu, y$ ⁴.

The classical Lagrangian is given by:

$$S = S_{gravity} + S_B , \quad (2)$$

$$S_{gravity} = \int d^4x dy \sqrt{-G} \{ -\Lambda(y) + \frac{1}{2} M_X^3 R \} , \quad (3)$$

$$S_B = \sum_{i=-l}^m \int d^4x \sqrt{-g^{(i)}} \{ \mathcal{L}_i - V_i \} , \quad (4)$$

where M_X is the 5-dimensional Planck scale, $\Lambda(y)$ is the cosmology constant, and V_i where $i = -l, \dots, m$ is the brane tension. The $\Lambda(y)$ is defined as the following:

$$\begin{aligned} \Lambda(y) = & \sum_{i=1}^m \Lambda_i (\theta(y - y_{i-1}) - \theta(y - y_i)) + \Lambda_{+\infty} \theta(y - y_m) \\ & + \sum_{i=-l+1}^0 \Lambda_i (\theta(-y + y_i) - \theta(-y + y_{i-1})) + \Lambda_{-\infty} \theta(-y + y_{-l}) , \end{aligned} \quad (5)$$

where $\theta(x) = 1$ for $x \geq 0$ and $\theta(x) = 0$ for $x < 0$. So, $\Lambda(y)$ is sectional constant.

The 5-dimensional Einstein equation for the above action is:

$$\begin{aligned} \sqrt{-G} \left(R_{AB} - \frac{1}{2} G_{AB} R \right) = & -\frac{1}{M_X^3} [\Lambda(y) \sqrt{-G} G_{AB} + \\ & \sum_{i=-l}^m V_i \sqrt{-g^{(i)}} g_{\mu\nu}^{(i)} \delta_M^\mu \delta_N^\nu \delta(y - y_i)] . \end{aligned} \quad (6)$$

Assuming that there exists a solution that respects 4-dimensional Poincare invariance in the x^μ -directions, one obtains the 5-dimensional metric:

$$ds^2 = e^{-2\sigma(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 . \quad (7)$$

⁴ We assume that $G_{\mu 5} = 0$ here. In addition, if the fifth dimesnion has Z_2 symmetry, i. e., the Lagrangian is invariant under the transformation $y \leftrightarrow -y$, then, $G_{\mu 5} = 0$.

With this metric, the Einstein equation reduces to:

$$\sigma'^2 = \frac{-\Lambda(y)}{6M_X^3} , \quad \sigma'' = \sum_{i=-l}^m \frac{V_i}{3M_X^3} \delta(y - y_i) . \quad (8)$$

The general solution to above differential equations is:

$$\sigma(y) = \sum_{i=-l}^m k_i |y - y_i| + k_c y + c , \quad (9)$$

where k_c and c are constants, and $k_i > 0$ for $i = -l, \dots, m$. The relations between the k_i and V_i , and the relations between the k_i and Λ_i are:

$$V_i = 6k_i M_X^3 , \quad \Lambda_i = -6M_X^3 \left(\sum_{j=i}^m k_j - \sum_{j=-l}^{i-1} k_j - k_c \right)^2 , \quad (10)$$

$$\Lambda_{-\infty} = -6M_X^3 \left(\sum_{j=-l}^m k_j - k_c \right)^2 , \quad \Lambda_{+\infty} = -6M_X^3 \left(\sum_{j=-l}^m k_j + k_c \right)^2 . \quad (11)$$

Therefore, the cosmology constant is negative except the section (at most one) with zero cosmology constant, then, for any point in $M^4 \times R^1$, which is not belong to any brane and the section with zero cosmology constant, there is a neighborhood which is diffeomorphic to (or a slice of) AdS_5 space. Moreover, the cosmology constant and brane tensions should satisfy above equations. In order to obtain finite 4-dimensional Planck scale, we obtain the constraints: $\sum_{j=-l}^m k_j > |k_c|$.

The general bulk metric is:

$$ds^2 = e^{-2\sum_{i=-l}^m k_i |y - y_i| - 2k_c y - 2c} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 . \quad (12)$$

And the corresponding 4-dimensional Planck scale is:

$$M_{pl}^2 = M_X^3 \left(T_{-\infty, -l} + T_{m, +\infty} + \sum_{i=-l}^{m-1} T_{i, i+1} \right) , \quad (13)$$

where

$$T_{-\infty, -l} = \frac{1}{2\chi_{-\infty}} e^{-2\sigma(y_{-l})} , \quad T_{m, +\infty} = \frac{1}{2\chi_{+\infty}} e^{-2\sigma(y_m)} , \quad (14)$$

if $\chi_{i, i+1} \neq 0$, then

$$T_{i, i+1} = \frac{1}{2\chi_{i, i+1}} \left(e^{-2\sigma(y_{i+1})} - e^{-2\sigma(y_i)} \right) , \quad (15)$$

and if $\chi_{i, i+1} = 0$, then

$$T_{i, i+1} = (y_{i+1} - y_i) e^{-2\sigma(y_i)} , \quad (16)$$

where

$$\chi_{\pm\infty} = \sum_{j=-l}^m k_j \pm k_c , \quad \chi_{i, i+1} = \sum_{j=i+1}^m k_j - \sum_{j=-l}^i k_j - k_c . \quad (17)$$

By the way, one can easily prove that $T_{i,i+1}$ is positive, which makes sure that the 4-dimensional Planck scale is positive.

In addition, the four dimensional GUT scale on i -th brane $M_{GUT}^{(i)}$ is related to the five dimensional GUT scale on i -th brane $M5_{GUT}^{(i)}$:

$$M_{GUT}^{(i)} = M5_{GUT}^{(i)} e^{-\sigma(y_i)} . \quad (18)$$

In this paper, we assume that $M5_{GUT}^{(i)} \equiv M_X$, for $i = -l, \dots, m$.

This solution can be generalized to the solution with Z_2 symmetry. Because of Z_2 symmetry, $k_c = 0$. There are two kinds of such models, one is the odd number of the branes, the other is the even number of the branes. For the first one, one just requires that $k_{-i} = k_i$, $y_{-i} = -y_i$, and $m = l$. For the second case, one just requires that $k_{-i} = k_i$, $y_{-i} = -y_i$, $m = l$, and $k_0 = 0$ (no number 0 brane). Furthermore, the solution can also be generalized to the case in which the fifth dimension is R^1/Z_2 , one just requires that $k_{-i} = k_i$, $y_{-i} = -y_i$, $m = l$, then, introduces the equivalence classes: $y \sim -y$ and i -th brane $\sim (-i)$ -th brane. The only trick point in this case is that the brane tension V_0 is half of the original value, i. e., $V_0 = 3k_0 M_X^3$.

3 Two Simple Models

Randall and Sundrum constructed the model with only one 3-brane which is the simplest model in above general construction [3]. However, the gauge hierarchy problem is not solved in that model. In this section, we will give two explicit simple models without gauge hierarchy problem, because all the other models with more branes will have these simple models as subset.

(I) Two brane cases with k_c : their positions are y_0, y_1 , respectively, and the values of the brane tension divided by $6M_X^3$ are: k_0, k_1 , respectively. And the constraint is $k_0 + k_1 > |k_c|$. Therefore, we obtain:

$$\sigma(y) = k_0|y - y_0| + k_1|y - y_1| + k_c y + c , \quad (19)$$

$$\sigma(y_0) = k_1(y_1 - y_0) + k_c y_0 + c , \quad \sigma(y_1) = k_0(y_1 - y_0) + k_c y_1 + c . \quad (20)$$

Assuming no section has zero cosmology constant, one obtains the four dimensional Planck scale:

$$M_{pl}^2 = M_X^3 \left(\frac{k_0}{k_0^2 - (k_1 - k_c)^2} e^{-2\sigma(y_0)} + \frac{k_1}{k_1^2 - (k_0 + k_c)^2} e^{-2\sigma(y_1)} \right) . \quad (21)$$

Without loss of generality, assuming that $\sigma(y_0) < \sigma(y_1)$ and $e^{-2(\sigma(y_1) - \sigma(y_0))} \ll 1$, we obtain

$$M_{pl}^2 = M_X^3 \frac{k_0}{k_0^2 - (k_1 - k_c)^2} e^{-2\sigma(y_0)} . \quad (22)$$

If the brane with position y_1 is the observable brane, assuming $M_X = \frac{k_0^2 - (k_1 - k_c)^2}{k_0}$, we obtain:

$$M_{GUT}^{(1)} = M_{pl} e^{-(\sigma(y_1) - \sigma(y_0))} . \quad (23)$$

So, we can push the GUT scale in our world to TeV scale and 10^5 GeV scale range if $(k_0 - k_1 + k_c)(y_1 - y_0) = 34.5$ and 30, respectively. And the value of $\sigma(y_0)$ determines the relation between M_{pl} and M_X . Explicit example: $k_0 = k_1 = k_c > 0$, $k_c(y_1 - y_0) = 34.5$ and 30. So, k_c is also an important factor to solve the gauge hierarchy problem.

If the brane with position y_0 is the observable brane, we can solve the gauge hierarchy problem only when $\sigma(y_0) > 0$. Assuming that $M_{pl} = \frac{k_0^2 - (k_1 - k_c)^2}{k_0}$ and $e^{-2(\sigma(y_1) - \sigma(y_0))} \ll 1$, we obtain:

$$M_{GUT}^{(0)} = M_{pl} e^{-\frac{1}{3}\sigma(y_0)} , \quad (24)$$

with $\sigma(y_0) = 103.5$ and 90, we can have GUT scale in our world at TeV scale and 10^5 GeV scale, respectively. The five-dimensional Planck scale is 10^{48} GeV and 10^{44} GeV, respectively.

(II) Three 3-branes with Z_2 symmetry, their positions are: $-y_1, 0, y_1$, respectively, and their brane tensions divided by $6M_X^3$ are: k_1, k_0, k_1 , respectively. One obtains M_{pl} , and the 4-dimensional GUT scales on the branes at $y=0$ and at $y=y_1$:

$$M_{pl}^2 = M_X^3 e^{-4k_1 y_1 - 2c} \left(\frac{1}{k_0} - \frac{2k_1}{(2k_1 + k_0)k_0} e^{-2k_0 y_1} \right) , \quad (25)$$

$$M_{GUT}^{(0)} = M5_{GUT}^{(0)} e^{-2k_1 y_1 - c} , \quad M_{GUT}^{(1)} = M5_{GUT}^{(1)} e^{-2k_1 y_1 - k_0 y_1 - c} , \quad (26)$$

where $M5_{GUT}^{(0)} = M5_{GUT}^{(1)} = M_X$. If the brane with position y_1 is the observable brane, one can solve the gauge hierarchy problem. For example, assuming $e^{-2k_0 y_1} \ll 1$ and $k_0 = M_X$, one obtains:

$$M_{pl} = M_X e^{-2k_1 y_1 - c} , \quad M_{GUT}^{(1)} = M_{pl} e^{-k_0 y_1} . \quad (27)$$

With $k_0 y_1 = 34.5$ and 30, one obtains the GUT scale in our world will be at TeV scale and 10^5 GeV scale range, respectively. And the relation between M_{pl} and M_X depends on the value of $2k_1 y_1 + c$. In addition, if the brane with position $y = 0$ is the observable brane, the gauge hierarchy problem can be solved, too. Assuming $M_{pl} = k_0$ and $e^{-2k_0 y_1}$ is very small, one obtains:

$$M_{pl} = M_X e^{-\frac{4}{3}k_1 y_1 - \frac{2}{3}c} , \quad M_{GUT}^{(0)} = M_{pl} e^{-\frac{2}{3}k_1 y_1 - \frac{1}{3}c} , \quad (28)$$

with $2k_1 y_1 + c = 103.5$ and 90, one can push the GUT scale in our world to the TeV scale and 10^5 GeV scale range, respectively. And the five dimensional Planck scale is 10^{48} GeV and 10^{44} GeV, respectively.

Acknowledgments

We would like to thank M. Cvetič, K. R. Dienes, C. Grojean, H. D. Kim and T. Shimizu for bringing my attention to their papers and helpful comments. This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation.

References

- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B**429** (1998) 263, hep-ph/9803315; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B**436** (1998) 257, hep-ph/9804398.
- [2] L. Randall and R. Sundrum, hep-ph/9905221.
- [3] L. Randall and R. Sundrum, hep-ph/9906064.
- [4] J. Lykken and L. Randall, hep-th/9908076.
- [5] M. Gogberashvili: hep-ph/9812296, hep-ph/9812365, hep-ph/9904383, hep-ph/9908347.
- [6] N. Arkani-Hamed, S. Dimopoulos, G. Dvali and N. Kaloper, hep-th/9907209.
- [7] W. D. Goldberger and M. B. Wise, hep-ph/9907447, hep-ph/9911457.
- [8] T. Nihei, hep-ph/9905487; N. Kaloper, hep-th/9905210.
- [9] C. Csaki, M. Grasesser, C. Kolda and T. Terning, hep-ph/9906513.
- [10] H. Verlinde, hep-th/9906182; A. Kehagias, hep-th/9908174.
- [11] I. Oda, hep-th/9908104.
- [12] A. Brandhuber and K. Sfetsos, hep-th/9908116.
- [13] H. B. Kim and H. D. Kim, hep-th/9909053.
- [14] T. Shiromizu, K. Maeda, M. Sasaki, gr-qc/9910076.
- [15] K. Behrndt and M. Cvetič, hep-th/9909058.
- [16] T. Li, hep-th/9908174, Phys. Lett. B to appear.
- [17] T. Li, hep-th/9912182.
- [18] T. Li, in preparation.
- [19] K. R. Dienes, E. Dudas, and T. Gherghetta, hep-ph/9908530.
- [20] O. DeWolfe, D.Z. Freedman, S.S. Gubser and A. Karch, hep-th/9909134.
- [21] S. Chang and M. Yamaguchi, hep-ph/9909523.
- [22] A. Chodos and E. Poppit, hep-th/9909199.
- [23] C. Grojean, J. Cline and G. Servant, hep-th/9906523, hep-ph/9909496, hep-th/9910081.
- [24] C. Csaki, M. Grasesser, L. Randall and T. Terning, hep-ph/9911406, and reference therein.
- [25] C. Csaki and Y. Shirman, hep-th/9908186; A. E. Nelson, hep-th/9909001.
- [26] I. Oda, hep-th/9909048.

- [27] H. Hatanaka, M. Sakamoto, M. Tachibana, and K. Takenaga, hep-th/9909076.
- [28] M. Cvetič and H. H. Soleng, Phys. Rept. **282** 159 (1997), hep-th/9604090, and references therein.
- [29] K.R. Dienes, E. Dudas, and T. Gherghetta, hep-ph/9803466, hep-ph/9806292, hep-ph/9807522.